

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09741 - Goat Anti-ACAT1 (aa253-266) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** ACAT1, acetyl-Coenzyme A acetyltransferase 1, ACAT, MAT, T2, THIL, acetoacetyl Coenzyme A thiolase, acetyl-CoA acetyltransferase 1, mitochondrial

acetoacetyl-CoA thiolase Official Symbol: ACAT1

Accession Number(s): NP\_000010.1

Human GeneID(s): 38

Non-Human GenelD(s): 110446 (mouse), 25014 (rat)

## **Immunogen**

Peptide with sequence C-DEEYKRVDFSKVPK, from the internal region of the protein sequence according to NP\_000010.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx. 40kDa band observed in lysates of cell lines CAC02, HEK293, HepG2, MCF7, NIH3T3 and KNRK and approx. 38-40kDa in Human, Mouse and Rat Liver lysates (calculated MW of 45.2kDa according to Human NP\_000010.1, 44.81kDa according to Mouse NP\_659033.1 and 44.7kDa according to Rat NP\_058771.2 ). Recommended concentration: 0.01-0.5μg/ml. Primary incubation 1 hour at room temperature.


**Immunofluorescence:** Strong expression of the protein seen in the cytoplasm of U2OS and A431 cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A431 cells. Recommended concentration: 10ug/ml.


# **Species Reactivity**

Tested: Human, Mouse, Rat


Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow




EB09741 (0.5μg/ml) staining of KNRK (A) and MCF7 (B), and (0.1μg/ml) HepG2 (C), Caco-2 (D), HEK293 (E), and NIH3T3 (F) cell lysate (35μg protein in RIPA buffer). Detected by chemiluminescence.



EB09741 (0.01μg/ml) staining of Human (A), and (0.1μg/ml) Mouse (B) and Rat (C), Liver lysate (35μg protein in RIPA buffer). Detected by chemilluminescence.



EB09741 Immunofluorescence analysis of paraformaldehyde fixed A431 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB09741 Flow cytometric analysis of paraformaldehyde fixed A431 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control:

Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.