

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07668 - Goat Anti-ACOX2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ACOX2, acyl-Coenzyme A oxidase 2, branched chain, BCOX, BRCACOX, BRCOX, THCCox, Peroxisomal branched chain acyl-CoA oxidase, THCA-CoA oxidase, Trihydroxycoprostanoyl-CoA oxidase

Official Symbol: ACOX2

Accession Number(s): NP_003491.1

Human GeneID(s): [8309](#)

Non-Human GeneID(s): 93732 (mouse), 252898 (rat)

Important Comments: Please note this antibody was designed using the mouse sequence, which differs by 1 amino acids from the human sequence.

Immunogen

Peptide with sequence C-HQSRLRPSPDPEAK, from the internal region (near the C Terminus) of the protein sequence according to NP_003491.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 75kDa band observed in Human Liver lysates (calculated MW of 76.8kDa according to NP_003491.1). Recommended concentration: 0.1-0.3µg/ml. An additional band of unknown identity was also consistently observed at 150kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Cow



EB07668 (0.1µg/ml) staining of Human Liver lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour.
Detected by chemiluminescence.