



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08207-B - Goat Anti-ACVR1, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ACVR1, activin A receptor type 1, ACTRI, ACVR1A, ACVRLK2, ALK2, FOP, SKR1, TSRI, TGF-B superfamily receptor type I, activin A receptor type I, activin A receptor, type I, activin A receptor, type II-like kinase 2, activin receptor type I, activin receptor-like kinase 2, hydroxyalkyl-protein kinase, serine/threonine-protein kinase receptor R1

Official Symbol: ACVR1

Accession Number(s): NP_001096.1

Human GeneID(s): [90](#)

Non-Human GeneID(s): 11477 (mouse), 79558 (rat)

Important Comments: Reported variants represent identical protein: NP_001096.1, NP_001104537.1

Immunogen

Peptide with sequence C-RKFKRRNQERLNPRD., from the internal region of the protein sequence according to NP_001096.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 60kDa band observed in Human Umbilical Cord lysates (calculated MW of 57.2kDa according to NP_001096.1). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



Biotinylated EB08207 (0.1µg/ml) staining of Human Umbilical Cord lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.