



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12488 - Goat Anti-Adipogenin (mouse, aa41-52) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: Adig, adipogenin, SMAF-1, Smaf1, small adipocyte factor 1

Official Symbol: Adig

Accession Number(s): NP_663610.1

Non-Human GeneID(s): 246747 (mouse)

Immunogen

Peptide with sequence SQESKEDDSDLC, from the internal region of the protein sequence according to NP_663610.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:2000.

Western blot: This antibody has been successfully used in WB on Mouse, PMID: 37249025.

IHC: Paraffin embedded Human Testis. Recommended concentration: 5µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Mouse

Specific Reference


This antibody has been successfully used in Western blot on Mouse:

Daisuke Aibara, Ai Sakaguchi, Kimihiko Matsusue

Transcriptional regulation of adipogenin expression in liver steatosis by hepatic peroxisome proliferator-activated receptor gamma

Genes Cells. 2023 Aug;28(8):585-594.

PMID: 37249025



EB12488 (5µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.