

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06937 - Goat Anti-ADRB2R / ADRB2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ADRB2, ADRB2R, adrenergic, beta-2-, receptor, surface [Homo sapiens], HGNC:286, ADRBR, B2AR, BAR, BETA2AR, beta-2 adrenergic receptor, beta-2 adrenoceptor, catecholamine receptor

Official Symbol: ADRB2

Accession Number(s): NP_000015.1

Human GeneID(s): [154](#)

Immunogen

Peptide with sequence C-HQGTVPSDNIDSQ, from the C Terminus of the protein sequence according to NP_000015.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 48kDa band observed in lysates of cell line HepG2 (calculated MW of 46.6kDa according to NP_000015.1). Recommended concentration: 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Dog, Cow

Specific Reference

This antibody (previous batch) has been successfully used in Western blot on Cow:
Rekawiecki R, Nowocin A, Kotwica J.

Relationship between concentrations of progesterone, oxytocin, noradrenaline, gene expression and protein level for their receptors in corpus luteum during estrous cycle in the cow.

Prostaglandins Other Lipid Mediat. 2010 Jun;92(1-4):13-8.

PMID: 20149890



EB06937 (0.5 μ g/ml) staining of HepG2 cell ysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.