



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06725 - Goat Anti-AGRP / Agouti-related protein Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** AGRP, ART, AGRT, ASIP2, agouti related protein homolog (mouse), agouti (mouse) related protein, Agouti-related transcript, mouse, homolog of, MGC118963, agouti related protein homolog

**Official Symbol:** AGRP

**Accession Number(s):** NP\_001129.1; NP\_015531.1

**Human GeneID(s):** [181](#)

**Important Comments:** This antibody is expected to recognise both reported isoforms (as represented by NP\_001129; NP\_015531)

### Immunogen

Peptide with sequence CRKLG TAMNPCSRT, from the C Terminus of the protein sequence according to NP\_001129.1; NP\_015531.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Preliminary experiments gave bands at approx 45kDa and 24kDa in Human Brain lysates after 0.1µg/ml antibody staining. These bands were successfully blocked by incubation with the immunizing peptide. Primary incubation 1 hour at room temperature. Please note that we cannot currently find an explanation in the literature given the calculated size of 14.4kDa according to NP\_001129.1 and NP\_015531.1.

**IHC:** Frozen section of Human Hypothalamus shows staining of a group of neuronal cell bodies in the human infundibular (arcuate) nucleus and their varicose axons.

Recommended concentration: 0.05-0.1µg/ml.

### Species Reactivity


**Tested:** Human

**Expected from sequence similarity:** Human, Dog, Pig

### Specific Reference

**This antibody has been successfully used as a specificity control in the following paper:**

Katalin Skrapits, Beáta Á Borsay, László Herczeg, Philippe Ciofi, Stephen R Bloom, Mohammad A Ghatei, Waljit S Dhillo, Zsolt Liposits, Erik Hrabovszky  
Colocalization of Cocaine- and Amphetamine-Regulated Transcript with Kisspeptin and Neurokinin B in the Human Infundibular Region  
PLoS One. 2014 Aug 1;9(8):e103977.  
PMID: 25084101



EB06725 (0.05µg/ml) staining of PFA-perfused cryosection of Human Hypothalamus. Antigen retrieval with citrate buffer Ph 6 at 80C for 30min, HRP-staining with Ni-DAB after Biotin-SP-antigoat amplification. Data obtained by Prof. Erik Hrabovszky, Inst, Exp, Med., Budapest, Hungary.