



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05419 - Goat Anti-AIF1/IBA1 isoform 1 and 3 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AIF1, IBA1, allograft inflammatory factor 1, AIF-1, IRT-1, interferon gamma responsive transcript, ionized calcium-binding adapter molecule 1, Daintain, DADB-70P7.8, allograft inflammatory factor-1 splice variant Hara-1, IRT1, protein G1

Official Symbol: AIF1

Accession Number(s): NP_116573.1; NP_001614.3

Human GeneID(s): [199](#)

Non-Human GeneID(s): 11629 (mouse), 29427 (rat)

Important Comments: This antibody is expected to recognize isoform 1 (NP_116573.1) and isoform 3 (NP_001614.3).

Immunogen

Peptide with sequence C-TGPPAKKAISELP, from the C Terminus of the protein sequence according to NP_116573.1; NP_001614.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 17kDa band observed in Mouse and Rat Brain lysates and in preliminary testing of Human Frontal Cortex lysate (calculated MW of 16.9kDa according to Mouse NP_062340.1 and 16.8kDa according to Rat NP_058892.1). Recommended concentration: 0.1-5µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Spleen and Brain (Frontal Cortex). Recommended concentration: 6-8µg/ml.

Species Reactivity

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Pig


Specific Reference

This antibody (previous batch) has been successfully used in IHC on Mouse:


Recio JS, Álvarez-Dolado M, Díaz D, Baltanás FC, Piquer-Gil M, Alonso JR, Weruaga E. Bone marrow contributes simultaneously to different neural types in the central nervous system through different mechanisms of plasticity.

Cell Transplant. 2011;20(8):1179-92.


PMID: 21294954



EB05419 (0.5 μ g/ml) staining of Mouse (A) and Rat (B) Brain lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.




EB05419 (6 μ g/ml) staining of paraffin embedded Human Spleen. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB05419 Negative Control showing staining of paraffin embedded Human Spleen, with no primary antibody.



EB05419 (6µg/ml) staining of paraffin embedded Human Frontal Cortex. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB05419 Negative Control showing staining of paraffin embedded Human Frontal Cortex, with no primary antibody.