



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05419-H - Goat Anti-AIF1/IBA1 isoform 1 and 3 (Human reactivity only) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** AIF1, IBA1, allograft inflammatory factor 1, AIF-1, IRT-1, interferon gamma responsive transcript, ionized calcium-binding adapter molecule 1, Daintain, DADB-70P7.8, allograft inflammatory factor-1 splice variant Hara-1, IRT1, protein G1

**Official Symbol:** AIF1

**Accession Number(s):** NP\_116573.1; NP\_001614.3

**Human GeneID(s):** [199](#)

**Important Comments:** This antibody is expected to recognize isoform 1 (NP\_116573.1) and isoform 3 (NP\_001614.3).

### Immunogen

Peptide with sequence C-TGPPAKKAISELP..., from the C Terminus of the protein sequence according to NP\_116573.1; NP\_001614.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx. 18kDa band observed in Human Frontal Cortex lysates (calculated MW of 16.7kDa according to NP\_001614.3). Recommended concentration: 1-3µg/ml.


Primary incubation 1 hour at room temperature.

**IHC:** Paraffin embedded Human Spleen. Recommended concentration: 4µg/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human



EB05419-H (2ug/ml) staining of Human Frontal Cortex lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB05419-H (4µg/ml) staining of paraffin embedded Human Spleen. Heat induced antigen retrieval with citrate buffer Ph 6, HRP-staining.



EB05419-H Negative Control showing staining of paraffin embedded Human Spleen, with no primary antibody.