



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05419-R - Goat Anti-AIF1/IBA1 isoform 1 and 3 (Rat reactivity only) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** AIF1, IBA1, allograft inflammatory factor 1, AIF-1, IRT-1, interferon gamma responsive transcript, ionized calcium-binding adapter molecule 1, Daintain, DADB-70P7.8, allograft inflammatory factor-1 splice variant Hara-1, IRT1, protein G1

**Official Symbol:** AIF1

**Accession Number(s):** NP\_116573.1; NP\_001614.3

**Non-Human GeneID(s):** 29427 (rat)

**Important Comments:** This antibody is expected to recognize isoform 1 (NP\_116573.1) and isoform 3 (NP\_001614.3).

### Immunogen

Peptide with sequence C-TGPPAKKAISELP., from the C Terminus of the protein sequence according to NP\_116573.1; NP\_001614.3.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 16kDa band observed in Rat Brain lysates (calculated MW of 16.8kDa according to Rat NP\_058892.1). Recommended concentration:1-3ug/ml. Primary incubation 1 hour at room temperature.

### Species Reactivity

**Tested:** Rat

**Expected from sequence similarity:** Rat



EB05419 (2.0 $\mu$ g/ml) staining of Rat Brain lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.