



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07257 - Goat Anti-AIF1 / IBA1 (isoforms 1 + 3) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: AIF1, IBA1, allograft inflammatory factor 1, DASS-82G15.2, AIF-1, IRT-1, Em:AF129756.17, interferon gamma responsive transcript, ionized calcium-binding adapter molecule

Official Symbol: AIF1

Accession Number(s): NP_116573.1; NP_001614.3

Human GeneID(s): [199](#)

Important Comments: This antibody is expected to recognise isoform 1 (NP_116573.1) and isoform 3 (NP_001614.3).

Immunogen

Peptide with sequence C-YEEKAREKEKP, from the C Terminus of the protein sequence according to NP_116573.1; NP_001614.3.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:16000.

IHC: In paraffin embedded Human Cerebellum shows staining of astrocytes.

Recommended concentration, 2-4µg/ml.


Flow Cytometry: Flow cytometric analysis of K562 cells. Recommended concentration: 10ug/ml.

Immunofluorescence: Strong expression of the protein seen in the cytoplasm of CaCo-2 cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human



EB07257 (4µg/ml) staining of paraffin embedded Human Cerebellum. Microwaved antigen retrieval with Tris/EDTA buffer pH9, HRP-staining.



EB07257 Flow cytometric analysis of paraformaldehyde fixed K562 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



EB07257 Immunofluorescence analysis of paraformaldehyde fixed Caco-2 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).