

#### International Office

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11511 - Goat Anti-AIMP1 / SCYE1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** AIMP1, aminoacyl tRNA synthase complex-interacting multifunctional protein 1, aminoacyl tRNA synthetase complex-interacting multifunctional protein 1, ARS-interacting multifunctional protein 1, EMAP2, EMAPII, endothelial monocyte-activating polypeptide 2, endothelial-monocyte activating polypeptide II, HLD3, multisynthase complex auxiliary component p43, multisynthetase complex auxiliary component p43, OTTHUMP00000161877, OTTHUMP00000161878, OTTHUMP00000219673, p43, SCYE1, small inducible cytokine subfamily E, member 1

(endothelial monocyte-activating)

Accession Number(s): NP\_004748.2; NP\_001135888.1

Human GeneID(s): 9255

Official Symbol: AIMP1

Non-Human GenelD(s): 13722 (mouse), 114632 (rat)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_004748.2; NP\_001135888.1). Reported variants represent identical protein:

NP\_004748.2, NP\_001135887.1

## Immunogen

Peptide with sequence NNDAVLKRLEQK-C, from the N Terminus of the protein sequence according to NP\_004748.2; NP\_001135888.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 37kDa band observed in lysates of cell lines HeLa and Jurkat, and in lysates of Mouse and Rat Spleen (calculated MW of 37kDa according to NP\_001135887.1). Recommended concentration: 0.3-1µg/ml.

### **Species Reactivity**

Tested: Human, Mouse, Rat

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow



EB11511 (0.3 $\mu$ g/ml) staining of Jurkat lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.