

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05538 - Goat Anti-Aldehyde Reductase Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** aldehyde reductase, AKR1A1, aldo-keto reductase family 1, member A1 (aldehyde reductase), ALDR1, ALR, MGC12529, MGC1380, alcohol dehydrogenase, aldo-keto reductase family 1, member A1, DD3, OTTHUMP00000009240, dihydrodiol dehydrogenase 3

Official Symbol: AKR1A1

Accession Number(s): NP\_006057.1; NP\_697021.1

Human GenelD(s): 10327

Non-Human GenelD(s): 58810 (mouse), 78959 (rat)

Important Comments: Both reported variants (NP\_006057.1and NP\_697021.1)

represent identical protein

# **Immunogen**

Peptide with sequence C-DAGHPLYPFNDPY, from the C Terminus of the protein sequence according to NP\_006057.1; NP\_697021.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.

**Western blot:** Approx 37kDa band observed in Human Liver lysates (calculated MW of 36.6kDa according to NP\_006057.1 and NP\_697021.1). Recommended concentration: 0.1-0.3µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


#### **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB05538 (0.1 $\mu$ g/ml) staining of Human Liver lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB05538 (5μg/ml) staining of paraffin embedded Human Kidney. Heat induced antigen retrieval with citrate buffer pH 6, HRP-staining.



EB05538 Negative Control showing staining of paraffin embedded Human Kidney, with no primary antibody.