

## **UK Office**

#### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB11052 - Goat Anti-Alk (mouse) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: ALK tyrosine kinase receptor, anaplastic lymphoma receptor tyrosine

kinase, CD246, CD246 antigen, NBLST3, Tcrz, Alk

Official Symbol: Alk

Accession Number(s): NP\_031465.2

Human GeneID(s): 238

Non-Human GenelD(s): 11682 (mouse), 266802 (rat)

## **Immunogen**

Peptide with sequence C-TETFHPERLESNS, from the internal region of the protein

sequence according to NP\_031465.2.

Please note the <u>peptide</u> is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum  $\,$ 

albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 150kDa band observed in Rat Brain lysates (calculated MW of

175kDa according to Rat NP\_001162572.1). Recommended concentration:

 $0.03-0.1\mu g/ml$ .

# **Species Reactivity**

Tested: Rat

Expected from sequence similarity: Mouse, Rat

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa

> 25kDa 20kDa

15kDa

EB11052 ( $0.03\mu g/ml$ ) staining of Rat Brain lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.