

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB06630 - Goat Anti-AMID / PRG3 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: AMID, PRG3, apoptosis-inducing factor (AIF)-like

mitochondrion-associated inducer of death, 5430437E11Rik, p53-responsive gene 3, apoptosis-inducing factor (AIF)-homologous mitochondrion-associated inducer of death, AIFM2, apoptosis indusing factor, mitochondrion associated 3, RB11, 367UE 2

AIFM2, apoptosis-inducing factor, mitochondrion-associated, 2, RP11-367H5.2,

OTTHUMP0000019740, apoptosis-inducing factor (AIF)-homologous

mitochondrion-associated inducer of death

Official Symbol: AIFM2

Accession Number(s): NP\_116186.1

Human GeneID(s): 84883

### **Immunogen**

Peptide with sequence GSQVSVESGALHC, from the N Terminus of the protein sequence according to NP\_116186.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Preliminary experiments gave an approx 48-50kDa band in Human Colon and Duodenum lysates after 0.1μg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 40.5kDa according to NP\_116186. The 48-50kDa band was successfully blocked by incubation with the immunizing peptide.


IHC: Paraffin embedded Human Skeletal Muscle and Heart. Recommended

concentration: 4µg/ml.


## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB06630 (4µg/ml) staining of paraffin embedded Human Skeletal Muscle. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB06630 (4μg/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.