

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07245-T - Goat Anti-Amphiphysin / AMPH Antibody - Trial

Size: 20µg specific antibody in 40µl



## **Target Protein**

Principal Names: AMPH, amphiphysin, amphiphysin (Stiff-Man syndrome with breast

cancer 128kDa autoantigen), HGNC:471, AMPH1

Official Symbol: AMPH

Accession Number(s): NP\_001626.1; NP\_647477.1

Human GeneID(s): 273

Non-Human GenelD(s): 218038 (mouse)

Important Comments: This antibody is expected to recognise both reported isoforms

(NP\_001626.1 and NP\_647477.1).

## **Immunogen**

Peptide with sequence C-KRKDESRISKAE, from the internal region of the protein sequence according to NP\_001626.1; NP\_647477.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 125kDa band observed in mouse brain lysates (calculated MW of 76.3kDa according to NP\_001626.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Wigge et al, Mol Biol Cell. 1997 Oct;8(10):2003-15.; PMID: 9348539). Recommended concentration: 0.1-0.5μg/ml. A minor band of unknown identity was also consistently observed at 45kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported? **IHC:** In paraffin embedded Human Cerebellum shows nuclear and fibrous staning. Recommended concentration, 2-4μg/ml.

## **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07245 ( $2.5\mu g/ml$ ) staining of paraffin embedded Human Cerebellum. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB07245 ( $0.5\mu g/ml$ ) staining of mouse brain lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.