

International Office

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07373 - Goat Anti-APBB1 / FE65 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: APBB1, FE65, amyloid beta (A4) precursor protein-binding, family B, member 1 (Fe65), MGC:9072, RIR, adaptor protein FE65a2, amyloid beta A4 precursor protein-binding, family B, member 1, stat-like protein
Official Symbol: APBB1
Accession Number(s): NP_001155.1; NP_663722.1
Human GenelD(s): 322
Important Comments: This antibody is expected to recognise both reported isoforms (NP_001155.1 and NP_663722.1).

Immunogen

Peptide with sequence C-GSLKPKRLGAHTP, from the C Terminus of the protein sequence according to NP_001155.1; NP_663722.1.

Please note the <u>peptide</u> is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 100kDa band observed in lysates of NIH/3T3 (calculated MW of 77.2kDa according to NP_001155.1 and 77.0kDa according to NP_663722.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Zambrano et al, J Biol Chem. 1998 Aug 7;273(32):20128-33.; PMID: 9685356). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

Species Reactivity

Tested: Mouse Expected from sequence similarity: Human

ł	250kDa 150kDa 100kDa 75kDa
	50kDa
	37kDa
	25kDa
	20kDa
	15kDa
	10kDa

EB07373 (0.1µg/ml) staining of NIH/3T3 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.