



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB07373 - Goat Anti-APBB1 / FE65 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** APBB1, FE65, amyloid beta (A4) precursor protein-binding, family B, member 1 (Fe65), MGC:9072, RIR, adaptor protein FE65a2, amyloid beta A4 precursor protein-binding, family B, member 1, stat-like protein

**Official Symbol:** APBB1

**Accession Number(s):** NP\_001155.1; NP\_663722.1

**Human GeneID(s):** [322](#)

**Important Comments:** This antibody is expected to recognise both reported isoforms (NP\_001155.1 and NP\_663722.1).

### Immunogen

Peptide with sequence C-GSLKPKRLGAHTP, from the C Terminus of the protein sequence according to NP\_001155.1; NP\_663722.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Approx 100kDa band observed in lysates of NIH/3T3 (calculated MW of 77.2kDa according to NP\_001155.1 and 77.0kDa according to NP\_663722.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Zambrano et al, J Biol Chem. 1998 Aug 7;273(32):20128-33.; PMID: 9685356). Recommended concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

### Species Reactivity

**Tested:** Mouse

**Expected from sequence similarity:** Human



EB07373 (0.1µg/ml) staining of NIH/3T3 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.