



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08357 - Goat Anti-APOBEC2 (aa12 - 26) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** APOBEC2, apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 2, ARCD1, ARP1, OTTHUMP00000039773, apolipoprotein B mRNA editing enzyme, catalytic polypeptide 2

**Official Symbol:** APOBEC2

**Accession Number(s):** NP\_006780.1

**Human GeneID(s):** [10930](#)

**Non-Human GeneID(s):** 11811 (mouse), 301226 (rat)

### Immunogen

Peptide with sequence C-EAASQNGEDLENLDD, from the internal region (near the N Terminus) of the protein sequence according to NP\_006780.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:1000.

**Western blot:** Approx 26kDa band observed in Human Heart and Skeletal Muscle lysates (calculated MW of 25.7kDa according to NP\_006780.1). Recommended concentration: 1-3µg/ml. An additional band of 90kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat



EB08357 (1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.