

### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08357 - Goat Anti-APOBEC2 (aa12 - 26) Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

**Principal Names:** APOBEC2, apolipoprotein B mRNA editing enzyme, catalytic polypeptide-like 2, ARCD1, ARP1, OTTHUMP00000039773, apolipoprotein B mRNA

editing enzyme, catalytic polypeptide 2

Official Symbol: APOBEC2

Accession Number(s): NP\_006780.1

Human GeneID(s): 10930

Non-Human GeneID(s): 11811 (mouse), 301226 (rat)

## **Immunogen**

Peptide with sequence C-EAASQNGEDLENLDD, from the internal region (near the N Terminus) of the protein sequence according to NP\_006780.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:1000.

**Western blot:** Approx 26kDa band observed in Human Heart and Skeletal Muscle lysates (calculated MW of 25.7kDa according to NP\_006780.1). Recommended concentration: 1-3µg/ml. An additional band of 90kDa was consistently observed, however this band was not blocked by the immunizing peptide and it is therefore a non-specific signal. We call for caution when used for other assays than Western blot.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat



EB08357 (1µg/ml) staining of Human Heart lysate (35µg protein in RIPA buffer) with (B) and without (A) blocking with the immunising peptide. Primary incubation was 1 hour. Detected by chemiluminescence.