



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB12282 - Goat Anti-APOD (aa70-80) Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** APOD, apolipoprotein D, apo-D

**Official Symbol:** APOD

**Accession Number(s):** NP\_001638.1

**Human GeneID(s):** [347](#)

### Immunogen

Peptide with sequence C-ENGKIKVLNQE, from the internal region of the protein sequence according to NP\_001638.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.


**Western blot:** Approx 26kDa band observed in Human Liver and Kidney lysates (calculated MW of 21.3kDa according to NP\_001638.1). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Adrenal Gland. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB12282 (0.3µg/ml) staining of Human Kidney (A) and Liver (B) lysates (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB12282 (5µg/ml) staining of paraffin embedded Human Adrenal Gland. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.