

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08329 - Goat Anti-Aromatase / CYP19A1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CYP19A1, cytochrome P450, family 19, subfamily A, polypeptide 1, ARO, ARO1, CPV1, CYAR, CYP19, MGC104309, P-450AROM, aromataseN: cytochrome P450, family 19, cytochrome P450, subfamily XIX (aromatization of androgens), estrogen

synthetase, flavoprotein-linked monooxygenase, microsomal monooxygenase

Official Symbol: CYP19A1

Accession Number(s): NP_000094.2; NP_112503.1

Human GeneID(s): 1588

Important Comments: This antibody is expected to recognise both reported isoforms

(NP_000094.2 and NP_112503.1), which are identical

Immunogen

Peptide with sequence C-HDLSLHPDETKN, from the internal region of the protein sequence according to NP_000094.2; NP_112503.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 50-55kDa band observed in Human Placenta lysates (calculated MW of 57.9kDa according to NP_000094.2 and NP_112503.1). Recommended


concentration: 0.1-0.3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Placenta. Recommended concentration: 4µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08329 (0.1 μ g/ml) staining of Human Placenta lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB08329 ($4\mu g/ml$) staining of paraffin embedded Human Placenta. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.