



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06745 - Goat Anti-Arrestin beta 2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** arrestin, beta 2, ARRB2, HGNC:712, ARB2, ARR2, DKFZp686L0365, arrestin beta 2, beta-arrestin 2, BARR2, arrestin 3, arrestin beta-2

**Official Symbol:** ARRB2

**Accession Number(s):** NP\_004304.1; NP\_945355.1; NP\_001244257.1; NP\_001244259.1; NP\_001244260.1; NP\_001316993.1

**Human GeneID(s):** [409](#)

**Non-Human GeneID(s):** 216869 (mouse), 25388 (rat)

**Important Comments:** This antibody is expected to recognise reported isoforms 1, 2, 3, 5, 6 and 7. No crossreactivity is expected with Arrestin beta 1.

### Immunogen

Peptide with sequence C-HDHIPLPRPQS, from the internal region of the protein sequence according to NP\_004304.1; NP\_945355.1; NP\_001244257.1; NP\_001244259.1; NP\_001244260.1; NP\_001316993.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.

**Western blot:** Approx 45kDa band observed in lysates of cell lines A549 and U251 and in preliminary testing of KNRK cell lysate (calculated MW of 46.1kDa according to NP\_004304.1 and 44.4kDa according to NP\_945355.1) Recommended concentration: 0.5-1µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in HeLa cells.

Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of A549 cells. Recommended concentration: 10ug/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat

### Specific References

**This antibody (previous batch) has been successfully used in IP on Rat:**

Zakrzewicz A, Krasteva G, Wilhelm J, Dietrich H, Wilker S, Padberg W, Wygrecka M, Grau V.

Reduced expression of arrestin beta 2 by graft monocytes during acute rejection of rat kidneys.

Immunobiology. 2011 Jul;216(7):854-61.

PMID: 21193245


**This antibody (previous batch) has been successfully used in WB on Mouse:**

Napolitano F, Pasqualetti M, Usiello A, Santini E, Pacini G, Sciamanna G, Errico F, Tassone A, Di Dato V, Martella G, Cuomo D, Fisone G, Bernardi G, Mandolesi G, Mercuri NB, Standaert DG, Pisani A.


Dopamine D2 receptor dysfunction is rescued by adenosine A2A receptor antagonism in a model of DYT1 dystonia.

Neurobiol Dis. 2010 Jun;38(3):434-45.


PMID: 20227500



EB06745 (1 $\mu$ g/ml) staining of A549 (A), and U251 (B) cell lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB06745 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing plasma membrane, cytoplasmic and nuclear staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06745 Flow cytometric analysis of paraformaldehyde fixed A549 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.