



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05291-B - Goat Anti-ASC / TMS1, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** PYCARD, PYD and CARD domain containing, ASC, CARD5, TMS, TMS-1, TMS1, caspase recruitment domain-containing protein 5, target of methylation-induced silencing 1

**Official Symbol:** PYCARD

**Accession Number(s):** NP\_037390.2; NP\_660183.1

**Human GeneID(s):** [29108](#)

**Important Comments:** This antibody is expected to recognise both reported isoforms (NP\_037390.2 and NP\_660183.1).

### Immunogen

Peptide with sequence CRESQSYLVEDLERS., from the C Terminus of the protein sequence according to NP\_037390.2; NP\_660183.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 24kDa band observed in lysates of cell line U937 (calculated MW of 21.6kDa according to NP\_037390.2). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



Biotinylated EB05291 (3 $\mu$ g/ml) staining of U937 lysate (35 $\mu$ g protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.