



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB12072 - Goat Anti-ASCL1 (aa79-91) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: ASCL1, achaete-scute complex homolog 1 (Drosophila), ASH1, HASH1, MASH1, bHLHa46, ASH-1, achaete scute protein, achaete-scute complex-like 1, achaete-scute homolog 1, class A basic helix-loop-helix protein 46

Official Symbol: ASCL1

Accession Number(s): NP_004307.2

Human GeneID(s): [429](#)

Non-Human GeneID(s): 17172 (mouse), 64186 (rat)

Immunogen

Peptide with sequence C-HKSAPKQVKRQRS, from the internal region of the protein sequence according to NP_004307.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Western blot: Approx 28kDa band observed in Mouse and Rat Lung lysates (calculated MW of 24.7kDa according to Mouse NP_032579.2 and 24.9kDa according to Rat NP_071779) . Recommended concentration: 0.3-1µg/ml.

Immunofluorescence: Strong expression of the protein seen in the nuclei and cytoplasm of MCF7 and U2OS cells. Recommended concentration: 10µg/ml.


Species Reactivity

Tested: Human, Mouse, Rat


Expected from sequence similarity: Human, Cow, Mouse, Rat, Dog



EB12072 (1 μ g/ml) staining of Mouse (A) and Rat (B) Lung lysate (35 μ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence



EB12072 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB12072 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nuclear and cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).