

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB12460 - Goat Anti-ATP5C1 (aa27-40) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** ATP5C1, ATP synthase, H+ transporting, mitochondrial F1 complex, gamma polypeptide 1, ATP5C, ATP5CL1, ATP synthase gamma chain, mitochondrial, ATP synthase subunit gamma, mitochondrial, F-ATPase gamma subunit, mitochondrial

ATP synthase, gamma subunit 1 **Official Symbol:** ATP5F1C

Accession Number(s): NP\_005165.1; NP\_001001973.1

Human GenelD(s): 509

Non-Human GenelD(s): 11949 (mouse), 116550 (rat)

**Important Comments:** The immunizing peptide represents the N terminus of the mature protein. This antibody is expected to recognize both reported isoforms (NP\_005165.1; NP\_001001973.1).

### **Immunogen**

Peptide with sequence C-TLKDITRRLKSIKN, from the internal region (near N terminus) of the protein sequence according to NP\_005165.1; NP\_001001973.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 30kDa band observed in Rat Skeletal Muscle lysates (calculated MW of 33.0kDa according to Rat NP\_446277.1). Recommended concentration: 0.1-0.3µg/ml.

### **Species Reactivity**

Tested: Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow



EB12460 (0.1 $\mu$ g/ml) staining of Rat Skeletal Muscle lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.