

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08038 - Goat Anti-B3GNT2 Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

Principal Names: B3GNT2, UDP-GlcNAc:betaGal

beta-1,3-N-acetylglucosaminyltransferase 2, B3GN-T1, B3GN-T2, B3GNT, B3GNT-2, B3GNT1, BETA3GNT, UDP-GlcNAc:betaGal beta-1,3-N-acetylglucosaminyltransferase 1,

beta-1,3-N-acetylglucosaminyltransferase bGnT-1, beta-1,3-N-a

Official Symbol: B3GNT2

Accession Number(s): NP\_006568.2

Human GeneID(s): 10678

Non-Human GeneID(s): 53625 (mouse)

### Immunogen

Peptide with sequence C-EKHKGFRTFDIE, from the internal region of the protein sequence according to NP\_006568.2.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 45kDa band observed in Human Pancreas lysates (calculated MW of 46.0kDa according to NP\_006568.2). Recommended concentration: 0.1-0.3µg/ml. An additional band of unknown identity was also consistently observed at 28kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field - have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat



EB08038 (0.1 $\mu$ g/ml) staining of Human Pancreas lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.