



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10369 - Goat Anti-BIRC3 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** AIP1, API2, apoptosis inhibitor 2, baculoviral IAP repeat-containing 3, baculoviral IAP repeat-containing protein 3, CIAP2, c-IAP2, HAIP1, HIAP1, MALT2, MIHC, RNF49, TNFR2-TRAF signaling complex protein, BIRC3

**Official Symbol:** BIRC3

**Accession Number(s):** NP\_001156.1

**Human GeneID(s):** [330](#)

**Important Comments:** Reported variants represent identical protein: NP\_892007.1, NP\_001156.1.

### Immunogen

Peptide with sequence C-SVNNLEATSQP, from the internal region of the protein sequence according to NP\_001156.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Experiments gave bands at approx 70kDa and 52kDa in Rat Testis lysates. These bands correspond to earlier findings with different antibodies from other sources.


This protein has a calculated MW of 67.3kDa according to Rat NP\_076477.2.

Recommended concentration: 1-3µg/ml.

### Species Reactivity

**Tested:** Rat

**Expected from sequence similarity:** Human



EB10369 (1 $\mu$ g/ml) staining of Rat Testis lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour.  
Detected by chemiluminescence.