

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08820 - Goat Anti-BMP2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: BMP2, bone morphogenetic protein receptor, type II (serine/threonine kinase), BMP2-IL, BMP2-III, BMP2, BRK-3, FLJ76945, PPH1, T-ALK, TRG10, bone morphogenetic protein receptor type II, primary pulmonary hypertension 1, serine/threonine kinase, transformation-related protein 10, type II activin receptor-like kinase

Official Symbol: BMP2

Accession Number(s): NP_001195.2

Human GeneID(s): [659](#)

Immunogen

Peptide with sequence C-DLETNKLDPKEVDK, from the internal of the protein sequence according to NP_001195.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 110kDa band observed in Human Liver lysates (calculated MW of 115kDa according to NP_001195.2). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Cow

Specific Reference

This antibody has been successfully used in the following paper:

James W. Connell, Rachel J. Allison, Catherine E. Rodger, Guy Pearson, Eliska Zlamalova, Evan Reid

ESCRT-III-associated proteins and spastin inhibit protrudin-dependent polarised membrane traffic.

Cell Mol Life Sci. 2020 Jul;77(13):2641-2658.

PMID: 31587092



EB08820 (1 μ g/ml) staining of Human Liver lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.