

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07979 - Goat Anti-BNIP1 Antibody

Size: 100µg specific antibody in 200µl



#### **Target Protein**

**Principal Names:** BNIP1, BCL2/adenovirus E1B 19kDa interacting protein 1, NIP1, SEC20, TRG-8, BCL2/adenovirus E1B 19kD interacting protein 1, BCL2/adenovirus E1B

19kD-interacting protein 1
Official Symbol: BNIP1

Accession Number(s): NP\_001196.1; NP\_053581.1; NP\_053582.1; NP\_053583.1

Human GeneID(s): 662

Non-Human GenelD(s): 224630 (mouse), 140932 (rat)

Important Comments: All reported variants represent isoform 1 (NP\_001196.1;

NP\_053581.1; NP\_053582.1; NP\_053583.1)

## **Immunogen**

Peptide with sequence C-DLLRQRKTTKESLAQ, from the internal region of the protein sequence according to NP\_001196.1; NP\_053581.1; NP\_053582.1; NP\_053583.1.

Please note the peptide is available for sale.

### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**


Peptide ELISA: antibody detection limit dilution 1:8000.

**Western blot:** Approx 28kDa band observed in Human Kidney lysates (calculated MW of 26.1kDa according to NP\_001196.1; NP\_053581.1; NP\_053582.1; NP\_053583.1). Recommended concentration: 0.5-1.5µg/ml. An additional band of unknown identity was also consistently observed at 48kDa. This band was successfully blocked by incubation with the immunising peptide. We would appreciate any feedback from people in the field have any such results been reported with other antibodies/lysates? Have any further splice variants/modified forms been reported?

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat



EB07979 (0.5 $\mu$ g/ml) staining of Human Kidney lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.