



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB09170 - Goat Anti-BRD4 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: BRD4, bromodomain containing 4, CAP, HUNK1, HUNKI, MCAP, bromodomain-containing 4, bromodomain-containing protein 4, chromosome-associated protein

Official Symbol: BRD4

Accession Number(s): NP_490597.1; NP_055114.1

Human GeneID(s): [23476](#)

Important Comments: This antibody is expected to recognize both reported isoforms (NP_490597.1; NP_055114.1).

Immunogen

Peptide with sequence C-PETSNPNKPKRQTNQ, from the internal region (near N Terminus) of the protein sequence according to NP_490597.1; NP_055114.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:8000.

Western blot: Approx 150kDa band observed in Mouse Testis lysates (calculated MW of 155.8kDa according to Mouse NP_065254.3). Recommended concentration: 0.5 -2µg/ml. Primary incubation was 1 hour.

Species Reactivity

Tested: Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB09170 (1 μ g/ml) staining of Mouse Testes lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.