



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05668 - Goat Anti-BRF2 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** BRF2, BRF2, subunit of RNA polymerase III transcription initiation factor, BRF1-like, BRFU, FLJ11052, TFIIIB50, transcription factor IIB- related factor, TFIIIB50, RNA polymerase III transcription initiation factor BRFU, RNA polymerase III transcription initiation factor BRF2

**Official Symbol:** BRF2

**Accession Number(s):** NP\_060780.2

**Human GeneID(s):** [55290](#)

### Immunogen

Peptide with sequence C-QAARQAATSVNPP, from the C Terminus of the protein sequence according to NP\_060780.2.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested


**Peptide ELISA:** antibody detection limit dilution 1:16000.

**Western blot:** Approx 50kDa band observed in HeLa lysates (predicted MW of 48kDa according to NP\_060780). Recommended for use at 1-3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human



EB05668 staining (1.5µg/ml) of HeLa lysate (RIPA buffer, 35µg total protein per lane). Primary incubated for 1 hour. Detected by chemiluminescence.