

#### **International Office**

Everest Biotech Ltd Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service: <u>customerservice@vectorlabs.com</u> Technical Service: <u>technical@vectorlabs.com</u>

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

## EB07531 - Goat Anti-CACNB4 (internal) Antibody

Size: 100µg specific antibody in 200µl

# **C**

#### **Target Protein**

Principal Names: CACNB4, calcium channel, voltage-dependent, beta 4 subunit, CAB4, CACNLB4, EA5, EJM, dihydropyridine-sensitive L-type, calcium channel beta-4 subunit, voltage dependent calcium channel beta 4 subunit Official Symbol: CACNB4 Accession Number(s): NP\_001005747.1; NP\_000717.2; NP\_001005746.1 Human GenelD(s): 785 Non-Human GenelD(s): 12298 (mouse), 58942 (rat) Important Comments: This antibody is expected to recognise all three reported isoforms (NP\_001005747.1; NP\_000717.2; NP\_001005746.1).

#### Immunogen

Peptide with sequence C-DYPDSYQDTYKPH, from the internal region (near the C Terminus) of the protein sequence according to NP\_001005747.1; NP\_000717.2; NP\_001005746.1.

Please note the peptide is available for sale.

#### **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.


#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Approx 55-60kDa band observed in Human Bone Marrow lysates (calculated MW of 54.7kDa according to NP\_001005747.1 and 58.2kDa according to NP\_000717.2). Recommended concentration:  $0.2-0.6 \ \mu g/ml$ .

### **Species Reactivity**

Tested: Human Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07531 (0.2µg/ml) staining of Human Bone Marrow lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.