

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08194 - Goat Anti-CADM4 (C Terminus) Antibody

Size: 100µg specific antibody in 200µl



# **Target Protein**

Principal Names: CADM4, cell adhesion molecule 4, IGSF4C, Necl-4, TSLL2, synCAM4,

TSLC1-like 2, immunoglobulin superfamily, member 4C, nectin-like 4

Official Symbol: CADM4

Accession Number(s): NP\_660339.1

Human GenelD(s): 199731

#### **Immunogen**

Peptide with sequence C-REAFLNGSDGHKRKE, from the C Terminus of the protein sequence according to NP\_660339.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.


**Western blot:** Approx 70kDa band observed in Human Brain (Cerebellum) lysates (calculated MW of 42.8kDa according to NP\_660339.1). The observed molecular weight corresponds to findings with EB08040, a product of different design reacting to a different part of the same protein. Recommended concentration: 0.03-0.1μg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Spleen and Liver. Recommended concentration: 2.5µg/ml.


#### **Species Reactivity**

Tested: Human


Expected from sequence similarity: Human, Mouse, Rat, Dog



Staining of Human Brain (Cerebellum) lysate (35 $\mu$ g protein in RIPA buffer). A) EB08040 (0.1 $\mu$ g/ml) and B) EB08194 (0.05 $\mu$ g/ml). Detected by chemiluminescence.



EB08194 (2.5 $\mu$ g/ml) staining of paraffin embedded Human Spleen. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB08194 (2.5μg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.