

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona,

CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07086 - Goat Anti-Calcipressin 1 / DSCR1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: DSCR1, Down syndrome critical region gene 1, HGNC:3040, ADAPT78, CSP1, DSC1, MCIP1, RCN1, RCAN1, Down syndrome candidate region 1, adapt78 protein, calcipressin 1, calcium and oxidant-inducible mRNA, modulatory calcineurin-interacting protein 1, myocyte-enriched calcineurin-interacting protein 1, near DSCR proline-rich protein, regulator of calcineurin 1

Official Symbol: DSCR1

Accession Number(s): NP_004405.3; NP_981962.1; NP_981963.1

Human GeneID(s): 1827

Non-Human GenelD(s): 54720 (mouse)

Important Comments: This antibody is expected to recognise all three reported isoforms

(NP_004405.3; NP_981962.1; NP_981963.1)

Immunogen

Peptide with sequence HVCESDQEKEE, from the internal region of the protein sequence according to NP_004405.3; NP_981962.1; NP_981963.1.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:1000.

Western blot: Western Blot: Preliminary experiments in Human Heart, Liver and Muscle lysates gave no specific signal but low background (at antibody concentration up to 1µg/ml). We would appreciate any feedback from people in the field - have any results been reported with other antibodies/lysates?

Species Reactivity

Tested

Expected from sequence similarity: Human, Mouse, Rat, Dog