

### **UK Office**

### **Everest Biotech Ltd**

Cherwell Innovation Centre

77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

Enquiries:

info@everestbiotech.com

Sales:

UK

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

#### **US Office**

### **Everest Biotech c/o Abcore**

405 Maple Street, Suite A106

Ramona, CA 92065 USA

Inquiries:

info@everestbiotech.com

Sales:

 $\underline{usasales@everest biotech.com}$ 

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09204 - Goat Anti-CAMK1D Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

Principal Names: CAMK1D, calcium/calmodulin-dependent protein kinase,

RP11-462F15.1, CKLiK, CaM-K1, CaMKID, CaM kinase ID, CamKI-like protein kinase,

OTTHUMP00000019119, OTTHUMP00000019120, OTTHUMP00000045030

Official Symbol: CAMK1D

Accession Number(s): NP\_065130.1

Human GeneID(s): 57118

Important Comments: This antibody is expected to recognize reported isoform 1

(NP\_065130.1).

## Immunogen

Peptide with sequence ASQKDCAYVAKPES, from the C Terminus of the protein sequence according to NP\_065130.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:4000.

**Western blot:** Approx. 40kDa band observed in Human Liver lysates (calculated MW of 40.2kDa according to NP\_065130.1). Recommended concentration: 1-3µg/ml.

### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human



EB09204 ( $0.5\mu g/ml$ ) staining of Human Liver lysate ( $35\mu g$  protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.