



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB11070 - Goat Anti-cardiolipin synthase 1 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** C20orf155, cardiolipin synthase, cardiolipin synthase 1, CLS, CLS1, dJ967N21.6, GCD10, CRLS1

**Official Symbol:** CRLS1

**Accession Number(s):** NP\_061968.1; NP\_001120930.1

**Human GeneID(s):** [54675](#)

**Important Comments:** This antibody is expected to recognize both reported isoforms (NP\_061968.1; NP\_001120930.1).

### Immunogen

Peptide with sequence C-RSALGSALDPLADK, from the internal region of the protein sequence according to NP\_061968.1; NP\_001120930.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.


**Western blot:** Approx 30kDa band observed in lysates of cell line HEK293 (calculated MW of 32.6kDa according to NP\_061968.1). Recommended concentration: 0.2-0.6µg/ml. Primary incubation was 1 hour.

**IHC:** Paraffin embedded Human Kidney. Recommended concentration: 5µg/ml.


### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow



EB11070 (0.2µg/ml) staining of HEK293 lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB11070 (5µg/ml) staining of paraffin embedded Human Kidney. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.