



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB08596 - Goat Anti-Cathepsin K Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CTSK, cathepsin K, CTS02, CTSO, CTSO1, CTSO2, MGC23107, PKND, PYCD, cathepsin K (pyncnodysostosis), cathepsin O1, cathepsin O2, cathepsin X

**Official Symbol:** CTSK

**Accession Number(s):** NP\_000387.1

**Human GeneID(s):** [1513](#)

### Immunogen

Peptide with sequence C-KTHRKQYNNKVDE, from the internal region (near N Terminus) of the protein sequence according to NP\_000387.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.


**Western blot:** Approx 37kDa band observed in Rat Skeletal Muscle lysates and in preliminary testing of Human Skeletal Muscle lysate (calculated MW of 36.8kDa according to Rat NP\_113748.1). Recommended concentration: 0.1-1µg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the vesicles of U2OS and MCF7 cells. Recommended concentration: 10µg/ml.


### Species Reactivity

**Tested:** Human, Rat


**Expected from sequence similarity:** Human, Mouse, Rat, Dog, Pig, Cow



EB08596 (1µg/ml) staining of Rat Skeletal Muscle lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB08596 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08596 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).