

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB08596 - Goat Anti-Cathepsin K Antibody

Size: 100µg specific antibody in 200µl



### **Target Protein**

**Principal Names:** CTSK, cathepsin K, CTS02, CTSO1, CTSO2, MGC23107, PKND, PYCD, cathepsin K (pycnodysostosis), cathepsin O1, cathepsin O2, cathepsin X

Official Symbol: CTSK

Accession Number(s): NP\_000387.1

Human GeneID(s): 1513

#### **Immunogen**

Peptide with sequence C-KTHRKQYNNKVDE, from the internal region (near N Terminus) of the protein sequence according to NP\_000387.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 37kDa band observed in Rat Skeletal Muscle lysates and in preliminary testing of Human Skeletal Muscle lysate (calculated MW of 36.8kDa according to Rat NP\_113748.1). Recommended concentration:0.1-1μg/ml. Primary incubation 1 hour at room temperature.

**Immunofluorescence:** Strong expression of the protein seen in the vesicles of U2OSand MCF7 cells. Recommended concentration:  $10\mu g/ml$ .


#### **Species Reactivity**

Tested: Human, Rat


Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB08596 (1µg/ml) staining of Rat Skeletal Muscle lysate (35µg protein in RIPA buffer). Detected by chemiluminescence.



EB08596 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB08596 Immunofluorescence analysis of paraformaldehyde fixed MCF7 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing vesicle staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).