



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06959-B - Goat Anti-CBX3 / HP1 Gamma, Biotinylated Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CBX3, chromobox 3, HECH, HP1-GAMMA, HP1Hs-gamma, HP1 gamma homolog, chromobox homolog 3 (HP1 gamma homolog, Drosophila), heterochromatin protein 1 homolog gamma, heterochromatin protein HP1 gamma, heterochromatin-like protein 1, modifier 2 protein

Official Symbol: CBX3

Accession Number(s): NP_009207.2

Human GeneID(s): [11335](#)

Non-Human GeneID(s): 12417 (mouse), 297093 (rat)

Important Comments: No cross-reactivity expected with HP1-alpha and HP1-beta.

Reported variants represent identical protein: NP_009207.2, NP_057671.2

Immunogen

Peptide with sequence CEAFLNSQKAGKEKD., from the internal region of the protein sequence according to NP_009207.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 20kDa band observed in lysates of cell line NIH3T3 (calculated MW of 20.8kDa according to Mouse NP_031650.3). See non-biotinylated parental product's datasheet for further QC data. Recommended concentration: 0.1-0.3µg/ml.

Species Reactivity

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Dog



Biotinylated EB06959 (0.3µg/ml) staining of NIH3T3 lysate (35µg protein in RIPA buffer), exactly mirroring its parental non-biotinylated product. Primary incubation was 1 hour. Detected by chemiluminescence, using streptavidin-HRP and using NAP blocker as a substitute for skimmed milk.