

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09667 - Goat Anti-CCL20 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** CCL20, chemokine (C-C motif) ligand 20, CKb4, LARC, MIP-3a, MIP3A, SCYA20, ST38, exodus-1, macrophage inflammatory protein 3 alpha, small

inducible cytokine subfamily A (Cys-Cys), member 20

Official Symbol: CCL20

Accession Number(s): NP\_004582.1; NP\_001123518.1

Human GeneID(s): 6364

**Important Comments:** This antibody is expected to recognize both isoforms

(NP\_004582.1; NP\_001123518.1).

## Immunogen

Peptide with sequence CLGYTDRILHPK, from the internal region of the protein sequence according to NP\_004582.1; NP\_001123518.1.

Please note the peptide is available for sale.

# **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin

Aliquot and store at -20°C. Minimize freezing and thawing.

# **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:16000.


**Western blot:** Preliminary experiments in Human Peripheral Blood Mononucleocytes lysates gave no specific signal but low background (at antibody concentration up to 2µg/ml).

**IHC:** In paraffin embedded Human Liver shows local staining between the hepatocytes. Recommended concentration, 6-8µg/ml.

## **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human, Cow, Pig



EB09667 (6μg/ml) staining of paraffin embedded Human Liver. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.