

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB09403 - Goat Anti-CCT3 / TCP1 Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** CCT3, chaperonin containing TCP1, subunit 3 (gamma), CCT-gamma, CCTG, PIG48, TCP-1-gamma, TRIC5, OTTHUMP00000025735, T-complex protein 1, gamma subunit, TCP1 (t-complex-1) ring complex, polypeptide 5, chaperonin containing TCP1, subunit 3

Official Symbol: CCT3

Accession Number(s): NP\_005989.3; NP\_001008883.1; NP\_001008800.1

Human GeneID(s): 7203

Non-Human GenelD(s): 12462 (mouse), 295230 (rat)

Important Comments: This antibody is expected to recognize all reported isoforms

(NP\_005989.3; NP\_001008883.1; NP\_001008800.1).

### **Immunogen**

Peptide with sequence C-GHKKKGDDQSRQGG, from the C-Terminus (near) of the protein sequence according to NP\_005989.3; NP\_001008883.1; NP\_001008800.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 65kDa band observed in Mouse Testis lysates (calculated MW of 60.5kDa according to Human NP\_005989.3 and 60.6kDa according to Mouse NP\_033966.1 and Rat NP\_954522.1). Recommended concentration: 0.01-0.03µg/ml.

IHC: Paraffin embedded Human Tonsil. Recommended concentration: 5µg/ml.


# **Species Reactivity**

Tested: Human, Mouse

Expected from sequence similarity: Human, Mouse, Rat, Pig, Cow, Dog



EB09403 (0.01µg/ml) staining of Mouse Testis lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.



EB09403 (5µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.