



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB05140 - Goat Anti-CD20 / MS4A1 (C Terminus) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: MS4A1, membrane-spanning 4-domains, subfamily A, member 1, B1, S7, BP35, CD20, LEU-16, MGC3969, CD20 antigen, CD20 receptor, B-lymphocyte cell-surface antigen B1, CVID5, leukocyte surface antigen Leu-16, MS4A2

Official Symbol: MS4A1

Accession Number(s): NP_068769.2; NP_690605.1

Human GeneID(s): [931](#)

Important Comments: Please note that both reported variants (NP_068769.2 and NP_690605.1) encode the same protein.

Immunogen

Peptide with sequence CQDQESSPIENDSSP, from the C Terminus of the protein sequence according to NP_068769.2; NP_690605.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested


Peptide ELISA: antibody detection limit dilution 1:16000.

Western blot: Approx 35kDa band observed in Human Lymph node lysates (calculated MW of 33.1kDa according to NP_068769.2). Recommended concentration: 0.05-0.3µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB05140 optimised QC. Primary incubation 1 hour at room temperature.
Image A: Human Lymph node lysate at primary Ab concentration 0.05ug/ml. (Loaded 35µg protein in RIPA buffer, per lane). Detected by chemiluminescence.