



## International Office

### Everest Biotech Ltd

Vector Laboratories, Inc.  
6737 Mowry Ave  
Newark, CA 94560  
United States

Customer Service:

[customerservice@vectorlabs.com](mailto:customerservice@vectorlabs.com)

Technical Service:

[technical@vectorlabs.com](mailto:technical@vectorlabs.com)

Tel: +1 (800) 227-6666

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB10740 - Goat Anti-CD28 Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** CD28, CD28 molecule, MGC138290, Tp44, CD28 antigen, CD28 antigen (Tp44), T-cell-specific surface glycoprotein

**Official Symbol:** CD28

**Accession Number(s):** NP\_006130.1

**Human GeneID(s):** [940](#)

**Important Comments:** The immunizing peptide corresponds to a part of the extracellular domain.

### Immunogen

Peptide with sequence SQQLQVYSKTGFNCD, from the internal region of the protein sequence according to NP\_006130.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:64000.


**Western blot:** Approx 48kDa band observed in Human Lymph Node and Tonsil lysates (calculated MW of 25.1kDa according to NP\_006130.1). The observed molecular weight corresponds to the glycosylated form. Recommended concentration: 0.3-1µg/ml. Primary incubation was 1 hour.

**IHC:** In paraffin embedded Human Lymph Node shows membrane staining in the paracortex. Recommended concentration: 1-2µg/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human



EB10740 (0.3 $\mu$ g/ml) staining of Human Tonsil lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.



EB10740 (2 $\mu$ g/ml) staining of paraffin embedded Human Lymph Node. Microwaved antigen retrieval with citrate buffer Ph 6, HRP-staining.



EB10740 Negative Control showing staining of paraffin embedded Human Lymph Node, with no primary antibody.