

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB07092 - Goat Anti-CDKN2A (isoform 3) Antibody

Size: 100µg specific antibody in 200µl



## **Target Protein**

**Principal Names:** CDKN2A, cyclin-dependent kinase inhibitor 2A (melanoma, p16, inhibits CDK4), HGNC:1787, ARF, CDK4I, CDKN2, CMM2, INK4, INK4a, MLM, MTS1, TP16, p14, p14ARF, p16, p16INK4, p16INK4a, p19, CDK4 inhibitor p16-INK4, cell cycle negative regulator beta, cyclin-dependent kinase inhibitor 2A, cyclin-dependent kinase inhibitor 2A, p16INK4a alternatively spliced form, cyclin-dependent kinase inhibitor p16, multiple tumor suppressor 1

Official Symbol: CDKN2A

Accession Number(s): NP\_478104.1

Human GeneID(s): 1029

Important Comments: This antibody is expected to recognize the pancreas isoform

(NP\_478104.1) only.

## **Immunogen**

Peptide with sequence C-ILRKKGRLPGEFSE, from the internal region of the protein sequence according to NP\_478104.1.

Please note the peptide is available for sale.

# Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

#### **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:128000.

**Western blot:** Approx 17kDa band observed in lysates of cell line HeLa (calculated MW of 12.1kDa according to NP\_478104.1). Recommended concentration: 1-3µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa 50kDa 37kDa 25kDa 20kDa

EB07091 (1 $\mu$ g/ml) staining of HeLa lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.