

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB07190-T - Goat Anti-CDX2 Antibody - Trial

Size: 20µg specific antibody in 40µl



Target Protein

Principal Names: CDX2, caudal type homeo box transcription factor 2, HGNC:1806, CDX-3, CDX3

Official Symbol: CDX2

Accession Number(s): NP_001256.1

Human GeneID(s): [1045](#)

Non-Human GeneID(s): 12591 (mouse), 66019 (rat)

Important Comments: No cross-reactivity expected with CDX1 and 4

Immunogen

Peptide with sequence EQLSPGGQRRNLCE, from the internal region of the protein sequence according to NP_001256.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:4000.

Western blot: Approx 35kDa band observed in Human Colon lysates (calculated MW of 33.5kDa according to NP_001256.1). Recommended concentration: 1-3µg/ml. Primary incubation was 1 hour.

IHC: Paraffin embedded Human Colon. Recommended concentration: 5µg/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog



EB07190 (1 μ g/ml) staining of Human Colon lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.



EB07190 (5 μ g/ml) staining of paraffin embedded Human Colon. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.