



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB06636 - Goat Anti-Choline acetyltransferase Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** choline acetyltransferase, CHAT, CMS1A, CMS1A2, acetyl  
CoA:choline O-acetyltransferase

**Official Symbol:** CHAT

**Accession Number(s):** NP\_066264.3; NP\_065574.3; NP\_001136405.1

**Human GeneID(s):** [1103](#)

**Important Comments:** This antibody is expected to recognise isoform 1 (NP\_066264.3), isoform 2 (NP\_065574.3) and isoform 3 (NP\_001136405.1). Reported variants represent identical protein (NP\_066264.3; NP\_066265.3; NP\_066266.3; NP\_001136406.1; NP\_001136401.1).

### Immunogen

Peptide with sequence CKEKATRPSQGHQP, from the C Terminus of the protein sequence according to NP\_066264.3; NP\_065574.3; NP\_001136405.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:32000.

**Western blot:** Approx 70kDa band observed in Human Placenta lysates (calculated MW of 70.5kDa according to NP\_065574 and NP\_066264). Recommended concentration: 0.03-0.3µg/ml.

### Species Reactivity

**Tested:** Human

**Expected from sequence similarity:** Human, Dog, Pig, Cow



EB06636 (0.03 $\mu$ g/ml) staining of Human Placenta lysate (35 $\mu$ g protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.