

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre 77 Heyford Park Upper Heyford Oxfordshire OX25 5HD

UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326 Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106 Ramona, CA 92065

USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB07335 - Goat Anti-CIAS1 / Cryopyrin (Internal) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CIAS1, cold autoinflammatory syndrome 1, AGTAVPRL, AII/AVP, C1orf7, FCAS, FCU, MWS, NALP3, PYPAF1, AII/AVP receptor-like, Muckle-Wells syndrome, NACHT, LRR and PYD containing protein 3, PYRIN-containing APAF1-like protein 1, angiotensin/vasopressin receptor AII/AVP-like, cryopyrin

Official Symbol: NLRP3

Accession Number(s): NP_004886.3; NP_899632.1; NP_001230062.1;

NP_001120933.1; NP_001120934.1

Human GeneID(s): 114548

Important Comments: This antibody is expected to recognise all five reported isoforms (NP_004886.3; NP_899632.1; NP_001230062.1; NP_001120933.1; NP_001120934.1). Reported variants represent identical protein (NP_004886.3 and NP_001073289.1).

Immunogen

Peptide with sequence C-DLYEKAKRDEPK, from the internal region of the protein sequence according to NP_004886.3; NP_899632.1; NP_001230062.1; NP_001120933.1; NP_001120934.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Preliminary experiments gave an approx 70kDa band in human lymph node and tonsil lysates after 0.3μg/ml antibody staining. Please note that currently we cannot find an explanation in the literature for the band we observe given the calculated size of 118kDa according to NP_004886.3 and 106kDa according to NP_899632.1. The 70kDa band was successfully blocked by incubation with the immunizing peptide.


IHC: In paraffin embedded Human Tonsil shows focal and interstitial staining of the

inflammatory elements close to the epithelia of the tonsil crypts. Recommended concentration: 2-4µg/ml.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat, Dog, Pig, Cow



EB07335 (2µg/ml) staining of paraffin embedded Human Tonsil. Steamed antigen retrieval with citrate buffer pH 6, HRP-staining.