

International Office

Everest Biotech Ltd

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

EB08550 - Goat Anti-CMG1 / CCDC2 / IFT74 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CMG1, CMG-1, CMG 1, Coiled-coil domain containing 2, Coiled-coil domain-containing protein 2, Capillary morphogenesis gene 1, Capillary morphogenesis protein 1, IFT74, intraflagellar transport 74 homolog, ntraflagellar transport protein 74 homolog

Official Symbol: IFT74

Accession Number(s): NP_001092692.1; NP_001092693.1; NP_079379.2; AAK77221.1

Human GeneID(s): 80173

Immunogen

Peptide with sequence CKTIVDALHSTSGN, from the C or N Terminus of the protein sequence according to NP_001092692.1; NP_001092693.1; NP_079379.2; AAK77221.1.

Please note the peptide is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.


Immunofluorescence: Strong expression of the protein seen in the cytoplasm and Golgi apparatus of U251 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of U251 cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human

Expected from sequence similarity: Human



EB08550 Immunofluorescence analysis of paraformaldehyde fixed U251 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml), showing cytoplasmic and Golgi apparatus staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (4ug/ml).



EB08550 Flow cytometric analysis of paraformaldehyde fixed U251 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (0.4ug/ml) showing weak staining. IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.