

#### **International Office**

#### **Everest Biotech Ltd**

Vector Laboratories, Inc. 6737 Mowry Ave Newark, CA 94560 United States

**Customer Service:** 

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

Research Use Only. Not for diagnostic or therapeutic use.

# EB05285-T - Goat Anti-COX1 / PTGS1 Antibody - Trial

Size: 20µg specific antibody in 40µl



## **Target Protein**

**Principal Names:** PTGS1, COX1, COX3, PHS1, PCOX1, PGHS1, PTGHS, PGG/HS, PGHS-1, prostaglandin-endoperoxide synthase 1 (prostaglandin G/H synthase and cyclooxygenase), prostaglandin G/H synthase and cyclooxygenase, RP11-542K23.6

Official Symbol: PTGS1

Accession Number(s): NP\_000953.2; NP\_542158.1

Human GenelD(s): 5742

Important Comments: This antibody is expected to recognise both human isoforms of

this protein according to NP\_000953.2 and NP\_542158.1.

## **Immunogen**

Peptide with sequence C-QDDGPAVERPSTEL, from the C Terminus of the protein sequence according to NP\_000953.2; NP\_542158.1.

Please note the peptide is available for sale.

## **Purification and Storage**

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

## **Applications Tested**

Peptide ELISA: antibody detection limit dilution 1:32000.

**Western blot:** Approx 70kDa band observed in lysates of histiocytic lymphoma cell line U937 (calculated MW of 68.7kDa according to NP\_000953.2) Recommended for use at 0.3-1µg/ml. Primary incubation 1 hour at room temperature.

**IHC:** In paraffin embedded Human Heart shows patchy staining of myocardial fibres in longitudinal section. Recommended concentration, 3-5µg/ml.

#### **Species Reactivity**

Tested: Human

Expected from sequence similarity: Human

250kDa 150kDa 100kDa 75kDa


37kDa

25kDa

20kDa

15kDa

EB05285 ( $1\mu g/ml$ ) staining of U937 cell lysate ( $35\mu g$  protein in RIPA buffer). Detected by chemiluminescence.



EB05285 (3.8 $\mu$ g/ml) staining of paraffin embedded Human Heart. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.