



International Office

Everest Biotech Ltd

Vector Laboratories, Inc.
6737 Mowry Ave
Newark, CA 94560
United States

Customer Service:

customerservice@vectorlabs.com

Technical Service:

technical@vectorlabs.com

Tel: +1 (800) 227-6666

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB08035 - Goat Anti-CRHR1 / CRF-R (aa 107 - 117) Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: CRHR1, CRF-R, corticotropin releasing hormone receptor 1, CRF1, CRFR1, CRH-R1h, CRHR, CRHR1f, corticotropin releasing hormone receptor variant 1h, seven transmembrane helix receptor

Official Symbol: CRHR1

Accession Number(s): NP_004373.2

Human GeneID(s): [1394](#)

Non-Human GeneID(s): 12921 (mouse), 58959 (rat)

Immunogen

Peptide with sequence C-NEEKKSQVHYH, from the internal region of the protein sequence according to NP_004373.2.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx. 55kDa band observed in Human Duodenum lysates and in preliminary testing of Human Ovary lysate (calculated MW of 47.7kDa according to NP_004373.2). This molecular weight is observed by other commercial sources, and was blocked by incubation with the immunizing peptide. A band at 48kDa was also observed in preliminary testing of Rat Testes lysate. Recommended concentration: 2-3µg/ml. Primary incubation 1 hour at room temperature.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human, Mouse, Rat

Specific References

This antibody (previous batch) has been successfully used in IHC on Mouse:

Cassandra L. Kooiker, Yuncai Chen, Matthew T. Birnie, and Tallie Z. Baram
Genetic Tagging Uncovers a Robust, Selective Activation of the Thalamic Paraventricular Nucleus by Adverse Experiences Early in Life.

Biol Psychiatry Glob Open Sci. 2023 Jan 24;3(4):746-755.

PMID: 37881549

This antibody (previous batch) has been successfully used in ICC on Mouse:

Silvia Cursano, Chiara R. Battaglia, Carolina Urrutia-Ruiz, Stefanie Grabrucker, Michael Schön, Jürgen Bockmann, Sonja Braumüller, Peter Radermacher, Francesco Roselli, Markus Huber-Lang, and Tobias M. Boeckers

A CRHR1 antagonist prevents synaptic loss and memory deficits in a trauma-induced delirium-like syndrome.

Mol Psychiatry. 2021; 26(8): 3778-3794.

PMID: 32051550

This antibody (previous batch) has been successfully used in ICC on Rat:

Xie Z, Penzes P, Srivastava DP

Exchange protein directly activated by cAMP 2 is required for corticotropin-releasing hormone-mediated

Eur J Neurosci. 2019 Jun 14.

PMID: 31199033

This antibody (previous batch) has been successfully used in ICC/IF on Rat:

Zhong Xie, Peter Penzes, Deepak P. Srivastava

EPAC2 is required for corticotropin-releasing hormone-mediated spine loss

April 2019: doi: <https://doi.org/10.1101/607598>

PMID: 0

This antibody (previous batch) has been successfully used in Western blot on Mouse:

Baglietto-Vargas D, Chen Y, Suh D, Ager RR, Rodriguez-Ortiz CJ, Medeiros R, Myczek K, Green KN, Baram TZ, LaFerla FM.

Short-term modern life-like stress exacerbates A β -pathology and synapse loss in 3xTg-AD mice.

J Neurochem. 2015 Sep;134(5):915-26.

PMID: 26077803

This antibody (previous batch) has been successfully used in Western blot on Mouse:

Weerachayaphorn J, Mennone A, Soroka CJ, Harry K, Hagey LR, Kensler TW, Boyer JL.

Nuclear factor-E2-related factor 2 is a major determinant of bile acid homeostasis in the liver and intestine.

Am J Physiol Gastrointest Liver Physiol. 2012 May 1;302(9):G925-36.

PMID: 22345550


This antibody (previous batch) has been successfully used in IHC on Mouse:

Jüngling K, Liu X, Lesting J, Coulon P, Sosulina L, Reinscheid RK, Pape HC.

Activation of neuropeptide S expressing neurons in the locus coeruleus by corticotropin-releasing factor.

J Physiol. 2012 May 8.

PMID: 22570383



EB08035 (2 μ g/ml) staining of Human Duodenum lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.